

NUCLEOTIDE SEQUENCES CODING FOR POLYPEPTIDES ENDOWED WITH A  
LARVICIDAL ACTIVITY TOWARDS LEPIDOPTERA

---

5           The subject of the invention is nucleotide sequences coding  
for polypeptides endowed with a larvicidal activity towards  
Lepidoptera.

10           It relates more particularly to agents, in particular  
nucleotide sequences, polypeptides or even vectors, or bacterial  
strains modified by these sequences and expressing polypeptides making  
it possible to prepare larvicidal compositions active against  
Lepidoptera, preferably against Spodoptera littoralis (hereafter  
S.littoralis) or Mamestra brassicae (hereafter designated by  
M.brassicae) or capable of transforming the plants to be treated in  
15           conferring on them this type of activity.

          It is known that most of the isolates of B.thuringiensis  
show a toxic activity with regard to larvae of more than a hundred  
species of Lepidoptera.

20           This activity results from the capacity of the strains of  
B.thuringiensis to synthesize, at the moment of sporulation,  
crystalline inclusions of protein nature, or  $\delta$ -endotoxins, under the  
control of one or several types of gene.

          It has been shown that the activity of these polypeptides  
is contained in the  $\text{NH}_2$ -terminal half or N-terminus of the protein.

25           The studies carried out have shown the high specificity  
of the  $\delta$ -endotoxins towards larvae of a given species.

          On account of this high specificity, many species of  
Lepidoptera, in particular of the family of the Noctuidae, react only  
weakly to commercial preparations of available B.thuringiensis.

30           It is so in particular for the species S.littoralis, a poly-  
phagous insect which constitutes the principal parasite of cotton  
and other industrially important crops. Among these crops, mention  
should be made of maize, the castor oil plant, tobacco, the groundnut,  
fodder plants, such as clover or alfalfa, or also market garden produce  
35           such as the cabbage or the tomato.

filed  
12/11/89  
6/5/95

Hence, one can imagine the interest of disposing of agents targeting specifically and effectively the family of the Noctuidae and in particular S.littoralis or M.brassicae.

5 The genes for  $\delta$ -endotoxins hitherto identified do not code for a polypeptide preferentially active with regard to S.littoralis.

The search by the inventors for a sequence of nucleotides coding for a polypeptide preferably active against the Noctuidae, more especially against S.littoralis, has led them to study the natural isolates of two strains of B.thuringiensis, the larvicidal activity of which on S.littoralis appears to be higher than that of the industrial preparations made starting from other strains of B.thuringiensis.

10 The species in question are aizawai 7-29 and entomocidus 6-01.

15 The study of these isolates has made it possible to demonstrate the existence of several genes for  $\delta$ -endotoxins of different structures and different specificities, of which two genes preferentially active against P.brassicae but not very active against the Noctuida of cotton and a gene inactive against P.brassicae and S.littoralis.

20 By studying the total DNA of these isolates and by carrying out appropriate hybridizations, followed by the cloning of the fragments identified by hybridization, the inventors have observed that it is possible to isolate nucleotide sequences implicated in genes for  $\delta$ -endotoxins coding for polypeptides active, preferably, against S.littoralis.

25 Thus, the aim of the invention is to provide nucleotide sequences capable of coding for at least the NH<sub>2</sub>-terminal part of a  $\delta$ -endotoxin toxic against the Noctuidae and preferably against S.littoralis or M.brassicae.

30 It also has the aim of providing a polypeptide toxic with regard to the Noctuidae.

35 Furthermore, the invention relates to a procedure for obtaining such a sequence and a polypeptide showing the desired activity as well as the intermediate agents such as vectors and

bacterial strains which can be utilized for obtaining the polypeptide.

In addition, the invention relates to the uses of these sequences and polypeptides for the development of larvicidal compositions with regard to the Noctuidae, in particular S.littoralis and for the transformation of the plants likely to be infected by these larvae.

The invention relates to a sequence of nucleotides coding for at least a part of the N-terminal region of a polypeptide toxic specifically against the larvae of Lepidoptera of the Noctuidae family, and preferably against S.littoralis, characterized by its capacity of hybridization with a gene capable of expressing a polypeptide toxic towards larvae of S.littoralis.

According to another aspect of the invention, the nucleotide sequence is characterized in that it is carried by a sequence of nucleotides of about 3 kb such as obtained by in vitro genetic recombination of sequences of nucleotides of B.thuringiensis capable of hybridizing with probes 1, 2 and 3 of pHTA2 shown in figure 2. The fragment of 3 kb corresponds more particularly to the restriction fragment HindIII-PstI.

The sequences of nucleotides of the invention are, in addition, characterized in that they contain sites in the following order : HindIII - HincII - BglIII - KpnI - HindIII - PstI.

In a preferred manner, these sequences of nucleotides are obtained by in vitro genetic recombination of DNA sequences derived from at least one strain of B.thuringiensis. In a variant of the embodiment of the invention, two different strains of B.thuringiensis are utilized.

Strains of B.thuringiensis particularly suited for obtaining these sequences of nucleotides are the strains corresponding to aizawai 7-29 and entomocidus 6-01, deposited on 21 April 1987 under the No. I-661 and No. I-660, respectively, with the National Collection of Cultures of Microorganisms (N.C.C.M.) in Paris.

In an advantageous manner, the sequences of nucleotides of the invention code for a polypeptide capable of forming an immunological complex with antibodies directed against polypeptides showing the larvicidal activity with regard to S.littoralis.



Sequences of nucleotides coding for at least a part of the N-terminal region of a polypeptide toxic specifically towards larvae of Lepidoptera of the Noctuidae family, and preferably towards S.littoralis, are characterized in that they contain the chain arrangement (I) defined above.

In an advantageous manner, the sequence of nucleotides characterized by the chain arrangement defined above codes for a part of a polypeptide having a higher larvicidal activity towards S.littoralis than that of the polypeptides encoded by natural isolates presently known for their effects against S.littoralis.

The study of this sequence of nucleotides shows that it is characterized by an initiation codon ATG situated at position 241 starting from which an open reading frame of 750 nucleotides has been identified.

This sequence is also characterized by a GGAGG attachment site for ribosomes at positions 230 to 234.

According to another feature, the sequence of nucleotides of the invention is characterized in that it contains, upstream from the ATG codon, a sequence going from the nucleotide at position 137 to the nucleotide at position 177, strongly homologous with the region found by Wong et al. (1983) and described in (16) upstream from the gene for the crystal of the strain kurstaki HD1 Dipel (BTK) and for which the authors have shown that it contains three promoters BtI, BtII and Ec which are functional in B.thuringiensis and E.coli, respectively. The homology of these sequences is about 70%.

The invention also relates to a sequence of nucleotides coding for the following sequence (II) of amino acids :

[illegible]

Author	Year	Country	Sample Size	Study Design	Findings
Wang et al.	2010	China	1,000	Case-control	Increased risk of lung cancer with high alcohol intake.
Li et al.	2011	China	2,000	Cohort	No significant association between alcohol and lung cancer.
Zhang et al.	2012	China	1,500	Case-control	Dose-response relationship between alcohol and lung cancer.
Chen et al.	2013	China	3,000	Cohort	Alcohol consumption associated with higher lung cancer risk.
Wu et al.	2014	China	1,200	Case-control	Significant association between alcohol and lung cancer.
Yang et al.	2015	China	2,500	Cohort	Alcohol intake linked to increased lung cancer incidence.
Xu et al.	2016	China	1,800	Case-control	High alcohol consumption increases lung cancer risk.
Guo et al.	2017	China	2,200	Cohort	Alcohol consumption associated with lung cancer.
Hou et al.	2018	China	1,600	Case-control	Alcohol intake significantly related to lung cancer.
Wang et al.	2019	China	2,800	Cohort	Alcohol consumption associated with lung cancer risk.
Li et al.	2020	China	1,400	Case-control	Alcohol intake associated with lung cancer.
Zhang et al.	2021	China	2,600	Cohort	Alcohol consumption associated with lung cancer.
Chen et al.	2022	China	1,900	Case-control	Alcohol intake associated with lung cancer.
Wu et al.	2023	China	2,100	Cohort	Alcohol consumption associated with lung cancer.
Yang et al.	2024	China	1,700	Case-control	Alcohol intake associated with lung cancer.
Xu et al.	2025	China	2,300	Cohort	Alcohol consumption associated with lung cancer.



TECHNICAL STAFF

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000



991  
 1001  
 1171  
 1261  
 1351  
 1441  
 1531  
 1621  
 1711  
 1801  
 1891  
 1981  
 2071  
 2161  
 2251  
 2341  
 2431  
 2521  
 2611  
 2701  
 2791  
 2881  
 2971  
 3061  
 3151  
 3241  
 3331  
 3421  
 3511  
 3601  
 3691  
 3781  
 3871  
 3961  
 4051  
 4141  
 4231  
 4321  
 4411  
 4501  
 4591  
 4681  
 4771  
 4861  
 4951  
 5041  
 5131  
 5221  
 5311  
 5401  
 5491  
 5581  
 5671  
 5761  
 5851  
 5941  
 6031  
 6121  
 6211  
 6301  
 6391  
 6481  
 6571  
 6661  
 6751  
 6841  
 6931  
 7021  
 7111  
 7201  
 7291  
 7381  
 7471  
 7561  
 7651  
 7741  
 7831  
 7921  
 8011  
 8101  
 8191  
 8281  
 8371  
 8461  
 8551  
 8641  
 8731  
 8821  
 8911  
 9001  
 9091  
 9181  
 9271  
 9361  
 9451  
 9541  
 9631  
 9721  
 9811  
 9901  
 9991  
 10081  
 10171  
 10261  
 10351  
 10441  
 10531  
 10621  
 10711  
 10801  
 10891  
 10981  
 11071  
 11161  
 11251  
 11341  
 11431  
 11521  
 11611  
 11701  
 11791  
 11881  
 11971  
 12061  
 12151  
 12241  
 12331  
 12421  
 12511  
 12601  
 12691  
 12781  
 12871  
 12961  
 13051  
 13141  
 13231  
 13321  
 13411  
 13501  
 13591  
 13681  
 13771  
 13861  
 13951  
 14041  
 14131  
 14221  
 14311  
 14401  
 14491  
 14581  
 14671  
 14761  
 14851  
 14941  
 15031  
 15121  
 15211  
 15301  
 15391  
 15481  
 15571  
 15661  
 15751  
 15841  
 15931  
 16021  
 16111  
 16201  
 16291  
 16381  
 16471  
 16561  
 16651  
 16741  
 16831  
 16921  
 17011  
 17101  
 17191  
 17281  
 17371  
 17461  
 17551  
 17641  
 17731  
 17821  
 17911  
 18001  
 18091  
 18181  
 18271  
 18361  
 18451  
 18541  
 18631  
 18721  
 18811  
 18901  
 18991  
 19081  
 19171  
 19261  
 19351  
 19441  
 19531  
 19621  
 19711  
 19801  
 19891  
 19981  
 20071  
 20161  
 20251  
 20341  
 20431  
 20521  
 20611  
 20701  
 20791  
 20881  
 20971  
 21061  
 21151  
 21241  
 21331  
 21421  
 21511  
 21601  
 21691  
 21781  
 21871  
 21961  
 22051  
 22141  
 22231  
 22321  
 22411  
 22501  
 22591  
 22681  
 22771  
 22861  
 22951  
 23041  
 23131  
 23221  
 23311  
 23401  
 23491  
 23581  
 23671  
 23761  
 23851  
 23941  
 24031  
 24121  
 24211  
 24301  
 24391  
 24481  
 24571  
 24661  
 24751  
 24841  
 24931  
 25021  
 25111  
 25201  
 25291  
 25381  
 25471  
 25561  
 25651  
 25741  
 25831  
 25921  
 26011  
 26101  
 26191  
 26281  
 26371  
 26461  
 26551  
 26641  
 26731  
 26821  
 26911  
 27001  
 27091  
 27181  
 27271  
 27361  
 27451  
 27541  
 27631  
 27721  
 27811  
 27901  
 27991  
 28081  
 28171  
 28261  
 28351  
 28441  
 28531  
 28621  
 28711  
 28801  
 28891  
 28981  
 29071  
 29161  
 29251  
 29341  
 29431  
 29521  
 29611  
 29701  
 29791  
 29881  
 29971  
 30061  
 30151  
 30241  
 30331  
 30421  
 30511  
 30601  
 30691  
 30781  
 30871  
 30961  
 31051  
 31141  
 31231  
 31321  
 31411  
 31501  
 31591  
 31681  
 31771  
 31861  
 31951  
 32041  
 32131  
 32221  
 32311  
 32401  
 32491  
 32581  
 32671  
 32761  
 32851  
 32941  
 33031  
 33121  
 33211  
 33301  
 33391  
 33481  
 33571  
 33661  
 33751  
 33841  
 33931  
 34021  
 34111  
 34201  
 34291  
 34381  
 34471  
 34561  
 34651  
 34741  
 34831  
 34921  
 35011  
 35101  
 35191  
 35281  
 35371  
 35461  
 35551  
 35641  
 35731  
 35821  
 35911  
 36001  
 36091  
 36181  
 36271  
 36361  
 36451  
 36541  
 36631  
 36721  
 36811  
 36901  
 36991  
 37081  
 37171  
 37261  
 37351  
 37441  
 37531  
 37621  
 37711  
 37801  
 37891  
 37981  
 38071  
 38161  
 38251  
 38341  
 38431  
 38521  
 38611  
 38701  
 38791  
 38881  
 38971  
 39061  
 39151  
 39241  
 39331  
 39421  
 39511  
 39601  
 39691  
 39781  
 39871  
 39961  
 40051  
 40141  
 40231  
 40321  
 40411  
 40501  
 40591  
 40681  
 40771  
 40861  
 40951  
 41041  
 41131  
 41221  
 41311  
 41401  
 41491  
 41581  
 41671  
 41761  
 41851  
 41941  
 42031  
 42121  
 42211  
 42301  
 42391  
 42481  
 42571  
 42661  
 42751  
 42841  
 42931  
 43021  
 43111  
 43201  
 43291  
 43381  
 43471  
 43561  
 43651  
 43741  
 43831  
 43921  
 44011  
 44101  
 44191  
 44281  
 44371  
 44461  
 44551  
 44641  
 44731  
 44821  
 44911  
 45001  
 45091  
 45181  
 45271  
 45361  
 45451  
 45541  
 45631  
 45721  
 45811  
 45901  
 45991  
 46081  
 46171  
 46261  
 46351  
 46441  
 46531  
 46621  
 46711  
 46801  
 46891  
 46981  
 47071  
 47161  
 47251  
 47341  
 47431  
 47521  
 47611  
 47701  
 47791  
 47881  
 47971  
 48061  
 48151  
 48241  
 48331  
 48421  
 48511  
 48601  
 48691  
 48781  
 48871  
 48961  
 49051  
 49141  
 49231  
 49321  
 49411  
 49501  
 49591  
 49681  
 49771  
 49861  
 49951  
 50041  
 50131  
 50221  
 50311  
 50401  
 50491  
 50581  
 50671  
 50761  
 50851  
 50941  
 51031  
 51121  
 51211  
 51301  
 51391  
 51481  
 51571  
 51661  
 51751  
 51841  
 51931  
 52021  
 52111  
 52201  
 52291  
 52381  
 52471  
 52561  
 52651  
 52741  
 52831  
 52921  
 53011  
 53101  
 53191  
 53281  
 53371  
 53461  
 53551  
 53641  
 53731  
 53821  
 53911  
 54001  
 54091  
 54181  
 54271  
 54361  
 54451  
 54541  
 54631  
 54721  
 54811  
 54901  
 54991  
 55081  
 55171  
 55261  
 55351  
 55441  
 55531  
 55621  
 55711  
 55801  
 55891  
 55981  
 56071  
 56161  
 56251  
 56341  
 56431  
 56521  
 56611  
 56701  
 56791  
 56881  
 56971  
 57061  
 57151  
 57241  
 57331  
 57421  
 57511  
 57601  
 57691  
 57781  
 57871  
 57961  
 58051  
 58141  
 58231  
 58321  
 58411  
 58501  
 58591  
 58681  
 58771  
 58861  
 58951  
 59041  
 59131  
 59221  
 59311  
 59401  
 59491  
 59581  
 59671  
 59761  
 59851  
 59941  
 60031  
 60121  
 60211  
 60301  
 60391  
 60481  
 60571  
 60661  
 60751  
 60841  
 60931  
 61021  
 61111  
 61201  
 61291  
 61381  
 61471  
 61561  
 61651  
 61741  
 61831  
 61921  
 62011  
 62101  
 62191  
 62281  
 62371  
 62461  
 62551  
 62641  
 62731  
 62821  
 62911  
 63001  
 63091  
 63181  
 63271  
 63361  
 63451  
 63541  
 63631  
 63721  
 63811  
 63901  
 63991  
 64081  
 64171  
 64261  
 64351  
 64441  
 64531  
 64621  
 64711  
 64801  
 64891  
 64981  
 65071  
 65161  
 65251  
 65341  
 65431  
 65521  
 65611  
 65701  
 65791  
 65881  
 65971  
 66061  
 66151  
 66241  
 66331  
 66421  
 66511  
 66601  
 66691  
 66781  
 66871  
 66961  
 67051  
 67141  
 67231  
 67321  
 67411  
 67501  
 67591  
 67681  
 67771  
 67861  
 67951  
 68041  
 68131  
 68221  
 68311  
 68401  
 68491  
 68581  
 68671  
 68761  
 68851  
 68941  
 69031  
 69121  
 69211  
 69301  
 69391  
 69481  
 69571  
 69661  
 69751  
 69841  
 69931  
 70021  
 70111  
 70201  
 70291  
 70381  
 70471  
 70561  
 70651  
 70741  
 70831  
 70921  
 71011  
 71101  
 71191  
 71281  
 71371  
 71461  
 71551  
 71641  
 71731  
 71821  
 71911  
 72001  
 72091  
 72181  
 72271  
 72361  
 72451  
 72541  
 72631  
 72721  
 72811  
 72901  
 72991  
 73081  
 73171  
 73261  
 73351  
 73441  
 73531  
 73621  
 73711  
 73801  
 73891  
 73981  
 74071  
 74161  
 74251  
 74341  
 74431  
 74521  
 74611  
 74701  
 74791  
 74881  
 74971  
 75061  
 75151  
 75241  
 75331  
 75421  
 75511  
 75601  
 75691  
 75781  
 75871  
 75961  
 76051  
 76141  
 76231  
 76321  
 76411  
 76501  
 76591  
 76681  
 76771  
 76861  
 76951  
 77041  
 77131  
 77221  
 77311  
 77401  
 77491  
 77581  
 77671  
 77761  
 77851  
 77941  
 78031  
 78121  
 78211  
 78301  
 78391  
 78481  
 78571  
 78661  
 78751  
 78841  
 78931  
 79021  
 79111  
 79201  
 79291  
 79381  
 79471  
 79561  
 79651  
 79741  
 79831  
 79921  
 80011  
 80101  
 80191  
 80281  
 80371  
 80461  
 80551  
 80641  
 80731  
 80821  
 80911  
 81001  
 81091  
 81181  
 81271  
 81361  
 81451  
 81541  
 81631  
 81721  
 81811  
 81901  
 81991  
 82081  
 82171  
 82261  
 82351  
 82441  
 82531  
 82621  
 82711  
 82801  
 82891  
 82981  
 83071  
 83161  
 83251  
 83341  
 83431  
 83521  
 83611  
 83701  
 83791  
 83881  
 83971  
 84061  
 84151  
 84241  
 84331  
 84421  
 84511  
 84601  
 84691  
 84781  
 84871  
 84961  
 85051  
 85141  
 85231  
 85321  
 85411  
 85501  
 85591  
 85681  
 85771  
 85861  
 85951  
 86041  
 86131  
 86221  
 86311  
 86401  
 86491  
 86581  
 86671  
 86761  
 86851  
 86941  
 87031  
 87121  
 87211  
 87301  
 87391  
 87481  
 87571  
 87661  
 87751  
 87841  
 87931  
 88021  
 88111  
 88201  
 88291  
 88381  
 88471  
 88561  
 88651  
 88741  
 88831  
 88921  
 89011  
 89101  
 89191  
 89281  
 89371  
 89461  
 89551  
 89641  
 89731  
 89821  
 89911  
 90001  
 90091  
 90181  
 90271  
 90361  
 90451  
 90541  
 90631  
 90721  
 90811  
 90901  
 90991  
 91081  
 91171  
 91261  
 91351  
 91441  
 91531  
 91621  
 91711  
 91801  
 91891  
 91981  
 92071  
 92161  
 92251  
 92341  
 92431  
 92521  
 92611  
 92701  
 92791  
 92881  
 92971  
 93061  
 93151  
 93241  
 93331  
 93421  
 93511  
 93601  
 93691  
 93781  
 93871  
 93961  
 94051  
 94141  
 94231  
 94321  
 94411  
 94501  
 94591  
 94681  
 94771  
 94861  
 94951  
 95041  
 95131  
 95221  
 95311  
 95401  
 95491  
 95581  
 95671  
 95761  
 95851  
 95941  
 96031  
 96121  
 96211  
 96301  
 96391  
 96481  
 96571  
 96661  
 96751  
 96841  
 96931  
 97021  
 97111  
 97201  
 97291  
 97381  
 97471  
 97561  
 97651  
 97741  
 97831  
 97921  
 98011  
 98101  
 98191  
 98281  
 98371  
 98461  
 98551  
 98641  
 98731  
 98821  
 98911  
 99001  
 99091  
 99181  
 99271  
 99361  
 99451  
 99541  
 99631  
 99721  
 99811  
 99901  
 10000

[illegible]

1591

1301  
 1302  
 1303  
 1304  
 1305  
 1306  
 1307  
 1308  
 1309  
 1310  
 1311  
 1312  
 1313  
 1314  
 1315  
 1316  
 1317  
 1318  
 1319  
 1320  
 1321  
 1322  
 1323  
 1324  
 1325  
 1326  
 1327  
 1328  
 1329  
 1330  
 1331  
 1332  
 1333  
 1334  
 1335  
 1336  
 1337  
 1338  
 1339  
 1340  
 1341  
 1342  
 1343  
 1344  
 1345  
 1346  
 1347  
 1348  
 1349  
 1350  
 1351  
 1352  
 1353  
 1354  
 1355  
 1356  
 1357  
 1358  
 1359  
 1360  
 1361  
 1362  
 1363  
 1364  
 1365  
 1366  
 1367  
 1368  
 1369  
 1370  
 1371  
 1372  
 1373  
 1374  
 1375  
 1376  
 1377  
 1378  
 1379  
 1380  
 1381  
 1382  
 1383  
 1384  
 1385  
 1386  
 1387  
 1388  
 1389  
 1390  
 1391  
 1392  
 1393  
 1394  
 1395  
 1396  
 1397  
 1398  
 1399  
 1400  
 1401  
 1402  
 1403  
 1404  
 1405  
 1406  
 1407  
 1408  
 1409  
 1410  
 1411  
 1412  
 1413  
 1414  
 1415  
 1416  
 1417  
 1418  
 1419  
 1420  
 1421  
 1422  
 1423  
 1424  
 1425  
 1426  
 1427  
 1428  
 1429  
 1430  
 1431  
 1432  
 1433  
 1434  
 1435  
 1436  
 1437  
 1438  
 1439  
 1440  
 1441  
 1442  
 1443  
 1444  
 1445  
 1446  
 1447  
 1448  
 1449  
 1450  
 1451  
 1452  
 1453  
 1454  
 1455  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465  
 1466  
 1467  
 1468  
 1469  
 1470  
 1471  
 1472  
 1473  
 1474  
 1475  
 1476  
 1477  
 1478  
 1479  
 1480  
 1481  
 1482  
 1483  
 1484  
 1485  
 1486  
 1487  
 1488  
 1489  
 1490  
 1491  
 1492  
 1493  
 1494  
 1495  
 1496  
 1497  
 1498  
 1499  
 1500  
 1501  
 1502  
 1503  
 1504  
 1505  
 1506  
 1507  
 1508  
 1509  
 1510  
 1511  
 1512  
 1513  
 1514  
 1515  
 1516  
 1517  
 1518  
 1519  
 1520  
 1521  
 1522  
 1523  
 1524  
 1525  
 1526  
 1527  
 1528  
 1529  
 1530  
 1531  
 1532  
 1533  
 1534  
 1535  
 1536  
 1537  
 1538  
 1539  
 1540  
 1541  
 1542  
 1543  
 1544  
 1545  
 1546  
 1547  
 1548  
 1549  
 1550  
 1551  
 1552  
 1553  
 1554  
 1555  
 1556  
 1557  
 1558  
 1559  
 1560  
 1561  
 1562  
 1563  
 1564  
 1565  
 1566  
 1567  
 1568  
 1569  
 1570  
 1571  
 1572  
 1573  
 1574  
 1575  
 1576  
 1577  
 1578  
 1579  
 1580  
 1581  
 1582  
 1583  
 1584  
 1585  
 1586  
 1587  
 1588  
 1589  
 1590  
 1591  
 1592  
 1593  
 1594  
 1595  
 1596  
 1597  
 1598  
 1599  
 1600  
 1601  
 1602  
 1603  
 1604  
 1605  
 1606  
 1607  
 1608  
 1609  
 1610  
 1611  
 1612  
 1613  
 1614  
 1615  
 1616  
 1617  
 1618  
 1619  
 1620  
 1621  
 1622  
 1623  
 1624  
 1625  
 1626  
 1627  
 1628  
 1629  
 1630  
 1631  
 1632  
 1633  
 1634  
 1635  
 1636  
 1637  
 1638  
 1639  
 1640  
 1641  
 1642  
 1643  
 1644  
 1645  
 1646  
 1647  
 1648  
 1649  
 1650  
 1651  
 1652  
 1653  
 1654  
 1655  
 1656  
 1657  
 1658  
 1659  
 1660  
 1661  
 1662  
 1663  
 1664  
 1665  
 1666  
 1667  
 1668  
 1669  
 1670  
 1671  
 1672  
 1673  
 1674  
 1675  
 1676  
 1677  
 1678  
 1679  
 1680  
 1681  
 1682  
 1683  
 1684  
 1685  
 1686  
 1687  
 1688  
 1689  
 1690  
 1691  
 1692  
 1693  
 1694  
 1695  
 1696  
 1697  
 1698  
 1699  
 1700  
 1701  
 1702  
 1703  
 1704  
 1705  
 1706  
 1707  
 1708  
 1709  
 1710  
 1711  
 1712  
 1713  
 1714  
 1715  
 1716  
 1717  
 1718  
 1719  
 1720  
 1721  
 1722  
 1723  
 1724  
 1725  
 1726  
 1727  
 1728  
 1729  
 1730  
 1731  
 1732  
 1733  
 1734  
 1735  
 1736  
 1737  
 1738  
 1739  
 1740  
 1741  
 1742  
 1743  
 1744  
 1745  
 1746  
 1747  
 1748  
 1749  
 1750  
 1751  
 1752  
 1753  
 1754  
 1755

...

[illegible]

•

LOW MTF AIR CTA GCA LAI GCA PCA TTY GAA BGA TCT CAT TTA GAA ABA GCA CAH ANG GCG GIG MAR' GLL CIG YIT ACT YCT YCC RAY  
2101

8

2131  
CAA AIC 666 TTA AAA ACC GAT GTG ACS GAT TAT CAT ATY CAA GTA TCC AAT TTA GTC GAT TGT CIG CAT

66A MAC-LLA 60A TT6 YCC 6AC RURA GTC RDA CAT 6CG RAG 6CA CTC RAS 6A1 6AG 666' RAR 11A C11 CDA 6A1 CCA MAC 11C RZA 666 ATC 2241

•

001 000 CAA CCA GAC CCG GGC AGA GGA AGT ACC ATC CAA GGA GGA GAT CAC GTA TTC AAA CAG AGT TAC ATC ACA CTA  
2431

11

CC6 C61 ACC 617 6A1 6A6 16C 1A1 CCA ACC 1A1 T1A TAI CAG AAA A1A 6A1 16C AAA T1A AAA 6C1 TAT ACC C61 TAT 6A4 11A A1A  
2321

;

CCG 1AT ATC GGA GAT AGT CAG GAC TTA GAG ATC 1AT TTG ATC GCG TAC AAT GCA AAA CAC GAG ATA GTA AAT GTG CCA GGC ACC GAT TCC  
2011

10

111A 106 CCG CTT 1CA GCG CAA AGT CCA ATC GGA AAG TGT GGA GAA CCG AAT CGA TCG GCG CCA CAC CTT GAA TGG AAT CCI GAT CTA GAT  
 2701

12

961 100 100 05

In a distinctive manner, sequences of nucleotides of the invention coding for a polypeptide toxic specifically towards larvae of Lepidoptera of the Noctuidae family, and preferably toward S.littoralis comprise or are constituted by the chain arrangement (III) previously defined.

The chain arrangement (III), comprised in the sequence of nucleotides of the invention contains 2711 nucleotides. This fragment includes in particular the potential promoter of the gene of the  $\delta$ -endotoxin active on S.littoralis.

Sequences of nucleotides modified in relation to the chain arrangements (I) or (III) described above naturally enter into the framework of the present invention to the extent to which these modifications do not generate appreciable variations of the toxicity of the polypeptide coded by the modified sequence towards S.littoralis.

These modifications may, for example, consist of deletions, substitutions, recombinations.

Thus, the sequences of nucleotides (I) and (III) contain at their position 611 a variable nucleotide corresponding to adenine (A) in the sequence (I) and to cytosine (C) in the sequence (III). These nucleotides enter into the composition of the respective codons GAA and GCA which code respectively for the amino acids glutamic acid (GLU) and alanine (ALA) in the respective sequences II and IV.

Similarly, any sequence of nucleotides which can hybridize with that of the chain arrangements (I) or (III) such as obtained by reverse enzymatic transformation of the corresponding RNA or even by chemical synthesis also enter into the framework of the definitions of the invention.

The sequence of nucleotides of formula (III) starts with a ATG initiation codon situated at position 241 and which represents the start of an open reading frame of 2470 nucleotides.

The invention also relates to a sequence of nucleotides characterized in that it codes for a polypeptide containing the sequence (IV) of amino acids below:

PET GLU GLU ASN ASN GLN ASN GLN CYS ILE

271

PRO TYR ASN CYS LEU SER ASN PRO GLU GLU VAL LEU LEU ASP GLY GLU ARG ILE SER THR GLY ASN SER ILE ASP ILE SER LEU SER

301

LEU VAL GLN PHE LEU VAL SER ASN PHE VAL PRO GLY GLY PHE LEU VAL GLY LEU ILE ASP PHE VAL TRP GLY ILE VAL GLY PRO SER

431

GLN TRP ASP ALA PHE LEU VAL GLN ILE GLU GLN LEU ILE ASN GLU ARG ILE ALA GLU PHE ALA ARG ASN ALA ALA ILE ALA ASN LEU GLU

541

GLY LEU GLY ASN ASN PHE ASN ILE TYR VAL GLU ALA PHE LYS GLU TRP GLU ASP PRO ASN ASN PRO ALA THR ARG THR ARG VAL ILE

631

ASP ARG PHE ARG ILE LEU ASP GLY LEU LEU GLU ARG ASP ILE PRO SER PHE ARG ILE SER GLY PHE GLU VAL PRO LEU LEU SER VAL TYR

721

ALA GLN ALA ALA ASN LEU HIS LEU ALA ILE LEU ARG ASP SER VAL ILE PHE GLY GLU ARG TRP GLY LEU THR THR ILE ASN VAL ASN GLU

811

ASN TYR ASN ARG LEU ILE ARG HIS ILE ASP GLU TYR ALA ASP HIS CYS ALA ASN THR TYR ASN ARG GLY LEU ASN ASN LEU PRO LYS SER

901

THR TYR GLN ASP TRP ILE THR TYR ASN ARG LEU ARG ARG ASP LEU THR THR VAL LEU ASP ILE ALA ALA PHE PHE PRO ASN TYR ASP

221

ASN ARG ARG TYR PRO ILE GLN PRO VAL GLY GLN LEU THR ARG GLU VAL TYR THR ASP PRO LEU ILE ASN PHE ASN PRO GLN LEU GLN SER

1061

VAL ALA GLN LEU PRO THR PHE ASN VAL MET GLU SER SER ALA ILE ARG ASN PRO HIS LEU PHE ASP ILE LEU ASN ASN LEU THR ILE PHE

1171

THR ASP TRP PHE SER VAL GLY ARG ASN PHE TYR TRP GLY GLY HIS ARG VAL ILE SER SER LEU ILE GLY GLY ASN ILE THR SER PRO

1261

ILE TYR GLY ARG GLU ALA ASN GLN GLU PRO PRO ARG SER PHE THR PHE ASN GLY PRO VAL PHE ARG THR LEU SER ILE PRO THR LEU ARG

1351

LEU LEU GLN GLN PRO CYS GLN ARG HIS PHE ASN LEU ARG GLY GLY VAL GLU PHE SER THR PRO THR ASN SER PHE THR TYR

1441

ARG GLY ARG GLY THR VAL ASP SER LEU THR GLU LEU PRO PRO GLU ASP ASN SER VAL PRO PRO ARG GLU GLY TYR SER HIS ARG LEU CYS

1531

HIS ALA THR PHE VAL GLN ARG SER GLY THR PRO PHE LEU THR THR GLY VAL PHE SER TRP THR HIS ARG SER ALA THR LEU THR ASN

1621

THR ILE ASP PRO GLU ARG ILE ASN GLN ILE PRO LEU VAL LYS GLY PHE ARG VAL TRP GLY GLY THR SER VAL ILE THR GLY PRO GLY PHE

1711

THR GLY GLY ASP ILE LEU ARG ARG ASN THR PHE GLY ASP PHE VAL SER LEU GLN VAL ASN ILE ASN SER PRO ILE THR GLN ARG TYR ARG

1801

LEU ARG PHE ARG TYR ALA SER SER ARG ASP ALA ARG VAL ILE VAL LEU THR GLY ALA SER THR GLY VAL GLY GLN VAL SER VAL

TTTGGG" SGGTGGG

1134

ASN PHE PRO LEU GLN LYS THR PHE GLU ILE GLY, GLU ASN LEU THR SER ARG THR PHE ARG TYR THR ASP PHE SER, ASN PRO PHE SER PHE

1261

ARG ALA ASN PRO ASP ILE ILE GLY ILE SER GLU GLN PRO LEU PHE GLY ALA GLY SER ILE SER SER GLY GLU LEU TYR ILE ASP LYS ILE

2071

GLU ILE ILE LEU ALA ASP ALA THR PHE GLU ALA, GLU SER ASP LEU GLU ARG ALA GLN LYS ALA VAL ASN ALA LEU PHE THR SER SER ASN

2161

GLN ILE GLY LEU LYS THR ASP VAL THR ASP TYR, HIS ILE ASP GLN VAL SER ASN LEU VAL ASP CYS LEU SER ASP GLU PHE CYS LEU ASP

2231

GLU LYS ARG GLU LEU SER GLU LYS VAL LYS HIS ALA LYS ARG LEU SER ASP, GLU ARG ASN LEU LEU GLN ASP PRO ASN PHE ARG GLY ILE

2241

ASN ARG GLN PRO ASP ARG GLY THR ARG GLY SER THR ASP ILE THR ILE GLN GLY GLY ASP VAL PHE LYS GLU ASN TYR VAL THR LEU

2431

PRO GLY THR VAL ASP GLU CYS TYR PRO THR TYR LEU TYR GLN LYS ILE ASP GLU SER LYS LEU LYS ALA THR THR ARG TYR GLU LEU ARG

2521

GLY TYR ILE GLU ASP SER GLN ASP LEU GLU ILE TYR LEU ILE ALA TYR ASN ALA LYS HIS GLU ILE VAL ASN VAL PRO GLY THR GLY SER

2611

LEU THR PRO LEU SER ALA GLN SER PRO ILE GLY LYS CYS GLY GLU PRO ASN ARG CYS ALA PRO HIS LEU GLU THR ASP PRO ASP LEU ASP

2761

CYS SER CYS

The invention also relates to recombinant expression and cloning vectors comprising more particularly at least one sequence of nucleotides such as that defined above, in particular at least a part of the sequence of about 3 kb.

5 A specific recombinant vector is, for example, a plasmid containing the HindIII-PstI fragment of the sequence of nucleotides of the invention, inserted in a vector pUC9. A first preferred vector is the plasmid pHT71, the construction of which is reported in the assemblies below, which comprises a HindIII-PstI DNA fragment according  
10 to the invention constituted uniquely of DNA derived from the strain aizawai 7-29.

Another recombinant vector is constituted by the plasmid pHT 671, the construction of which is given in figure 4. This plasmid contains a chimeric HindIII-PstI fragment, obtained by fusing a HindIII-HindII fragment of 1.1 kb derived from the strain entomocidus 6-01  
15 with a HincII-PstI fragment of 1.9 kb derived from the strain aizawai 7-29.

The modified bacterial strains which contain one of the nucleotide sequences defined above or also a recombinant expression  
20 vector and cloning previously defined, and preferably the plasmid pHT671 or the plasmid pHT71, also enter into the framework of the invention.

The invention also relates to a polypeptide toxic towards larvae of Lepidoptera and in a preferential manner towards  
25 S.littoralis, which attack cotton leaves or other crops such as those listed above, characterized in that it is capable of forming an immunological complex with antibodies directed against polypeptides with larvicidal activity towards S.littoralis.

The invention relates more particularly to the NH<sub>2</sub>-terminal  
30 part of this polypeptide which contains the larvicidal activity.

The extremity of the active NH<sub>2</sub>-terminal part corresponds to the sequence (II) of amino acids given above.

A preferred polypeptide of the invention is that which corresponds to this sequence (II) and corresponds to the sequence  
35 (IV) of amino acids given in the preceding pages. This polypeptide corresponding to the sequence (IV) contains 823 amino acids. Its calculated molecular mass is 92906 Da.

This sequence of  $\delta$ -endotoxin was compared with amino acid sequences of  $\delta$ -endotoxins derived from other strains of B.thuringiensis active on the Lepidoptera and the genes of which have been isolated and sequenced previously : the  $\delta$ -endotoxins in question are those of the strains kurstaki HD1 (19), kurstaki HD73 (20), berliner 1715 (21) and (22) Sotto (23) and aizawai IPL7 (24).

The results of these comparisons indicate that all are different in the second quarter of the molecule (amino acids 281 to 620) whereas the  $\text{NH}_2$ -terminal part (amino acids 1 to 280) and the COOH-terminal domain (amino acids 621 to 1175) of the protein are highly conserved and differ only by several amino acids. On the other hand, the  $\delta$ -endotoxin corresponding to the sequence (IV) above shows considerable differences from the other  $\delta$ -endotoxins both in the  $\text{NH}_2$ -terminal part (amino acids 1 to 280) and in the second quarter of the molecule (amino acids 281 to 620). The results of these comparisons indicate again that the  $\text{NH}_2$ -terminal half of the molecule (amino acids 1 to 620) which corresponds to the toxic fraction of the protein only show 46% homology with the other  $\delta$ -endotoxins. The most important differences are located in the second half of the toxic part of the molecule (amino acids 281 to 620) with only 36% of identical amino acids, the  $\text{NH}_2$ -terminal part (amino acids 1 to 280) itself showing 58% of amino acids identical with the other  $\delta$ -endotoxins. Such considerable differences have never been observed up to now in the  $\text{NH}_2$ -terminal part of the toxic fraction of the molecule among the  $\delta$ -endotoxins active on the Lepidoptera.

In order to obtain a sequence of nucleotides entering into the framework of the invention, coding for at least the active part of a polypeptide showing a specific toxicity towards larvae of Lepidoptera of the Noctuidae family, and preferably towards S.littoralis, recourse is had, in conformity with the invention, to the following steps, namely:

- the carrying out of a molecular hybridization between, on the one hand, a nucleotide sequence of a strain of B.thuringiensis active against S.littoralis and, on the other, at least two nucleotide sequences, used as probes, derived from the 5' part of a restriction



fragment of a gene for  $\delta$ -endotoxin of B.thuringiensis, this part coding for the NH<sub>2</sub>-terminal part of the polypeptide active on the larvae of Lepidoptera, and from the 3' part of this fragment coding for the COOH part of the polypeptide,

- 5 - the isolation of the hybrid fragment,
- its cloning in a vector, followed by its purification.

In an advantageous manner, the hybridization probes utilized are obtained from a gene for the  $\delta$ -endotoxin derived from the strain aizawai 7-29 coding for a protein of 130 kDa, active against P.brassicae and inactive towards S.littoralis, this gene having been  
10 cloned in the recombinant plasmid pHTA2.

In an embodiment of the preceding procedure, the fragment recombined with the vector in the cloning step is elaborated from a HindIII-PstI restriction fragment derived from a single strain of  
15 B.thuringiensis, preferably aizawai 7-29. In particular, this fragment is carried preferentially by the recombinant plasmid pHTA6 such as isolated with the aid of a probe constituted by a PvuII fragment of 2 kb of the plasmid pBT15-88 corresponding to the internal part of a gene for the chromosomal crystal of the strain berliner 1715,  
20 starting from transforming clones containing nucleotide sequences derived from B.thuringiensis strains active against larvae of Lepidoptera, inter-alia of S.littoralis.

In another embodiment of the invention, the fragment recombined with the vector in the cloning step is elaborated from  
25 several sequences of nucleotides derived from recombinant vectors containing sequences of nucleotides from at least two different strains of B.thuringiensis, possessing the same restriction maps and themselves containing all or part of the sequences of nucleotides capable of coding for a polypeptide active, in a preferential manner, against  
30 S.littoralis.

In this case, the recombined fragment used in the cloning step is a fragment of about 3 kb, advantageously elaborated from a HindIII-HincII restriction fragment of about 1.1 kb derived from the entomocidus 6-01 strain and a HincII-PstI fragment of about 1.9 kb  
35 from the aizawai 7-29 strain. It corresponds to a truncated gene for  $\delta$ -endotoxin.

The HindIII-HincII and HincII-PstI restriction fragments are carried more especially by the respective recombinant plasmids pHTE6 and pHTA6 such as isolated with the aid of the probe constituted by the PvuII fragment mentioned above.

5           The study of the toxicity towards the larvae of Lepidoptera of the bacterial strains modified with the aid of the sequences of nucleotides defined above, has made it possible to demonstrate their high toxic activity, in particular with regard to the caterpillars of S.littoralis.

10           This activity was estimated from the point of view of the specificity index corresponding to the ratio

LC50 S.littoralis

---

LC50 P.brassicae

15           in which "LC50" represents the lethal concentration killing 50% of the larvae in 72 hours.

          The utilization of such an index makes it possible to evaluate the activity of the bacterial strains studied without having to consider the level of expression of the polypeptides.

20           The results obtained, which are reported in the examples which follow, and the values of LD50 which are deduced from them, have shown that the bacterial strains modified according to the invention show a more specific toxic activity towards S.littoralis than the native crystal proteins of the strains aizawai 7-29 or berliner 1715.

25           Therefore, the invention relates to the use of these modified strains, of recombinant vectors containing the nucleotide sequences defined above, in particular the plasmid pHT671 and the plasmid pHT71, and these sequences themselves for the elaboration of larvicidal compositions preferentially toxic towards S.littoralis.

30           The larvicidal compositions of the invention are thus characterized in that they contain an efficacious quantity of polypeptides such as defined above or expressed by the nucleotide sequences mentioned above.

35           In order to produce these polypeptides the truncated genes

for  $\delta$ -endotoxin corresponding to the nucleotide sequences of the invention are advantageously implemented.

These genes can be used to produce in excess the toxic polypeptide in microorganisms permitting the expression of the above recombinant vectors. Suitable strains of microorganisms include E.coli or also B.subtilis.

These truncated genes may be reintroduced into the strains of B.thuringiensis or into related species such as B.cereus, according to the standard techniques, for example, by transformation, conjugation or transduction. These techniques make it possible to produce the toxic polypeptide in large quantity without first having to modify the natural region of the promoter for the  $\delta$ -endotoxin genes of B.thuringiensis.

This transformation may be carried out by using methods derived from the transformation of the protoplasts of B.subtilis according to (11) or of the vegetative cells of B.thuringiensis as described in (12).

The introduction of recombinant plasmids by a system of the conjugation type may be carried out by using B.thuringiensis as host strain and B.subtilis or Streptococcus faecalis as strains of the donor type by operating according to (13) and (14).

As a variant, the sequences of nucleotides are introduced into microorganisms living in the environment or in association with the plants and capable of expressing recombinant vectors containing these sequences. The introduction may be carried out in microorganisms such as Pseudomonas by working according to the procedure described in (17), by the intermediary of plasmid vectors containing the transposon Tn5 and the gene for the toxin, or Azospirillum or Rhizobium by means of the intermediary of suicide vectors derived from the plasmid RP4 and of mobilizing plasmids functional in Gram negative bacteria (for example, pRK2013) according to the procedures described in (18).

The truncated genes are alone in the strains of Bacilli or, as a variant, are associated with different  $\delta$ -endotoxin genes which makes it possible to obtain crystals synthesized by these strains

specifically toxic towards given species of Noctuidae, or toxic both towards the Noctuidae and insects sensitive to other  $\delta$ -endotoxins. These recombinations, carried out in vitro or in vivo with the nucleotide sequences of the invention and other  $\delta$ -endotoxin genes showing different toxic specificities, lead to the construction of new genes coding for novel hybrid toxic proteins exhibiting a large spectrum of activity towards insects. These new genes and these novel proteins also enter into the framework of the invention.

In these applications, the nucleotide sequences of the invention may be transferred and expressed in plants sensitive to S.littoralis in order to diminish the devastation caused by these insects.

Among the plants to be protected, mention should be made of : cotton, clover, the tomatoe and alfalfa.

The transfer of the truncated gene into cotton plants may be carried out by transformation involving strains such as Agrobacterium as described in (15).

In addition, the invention relates to the plant cells, the plants and the seeds containing the nucleotide sequences defined above.

The plant cells according to the invention are cells, the genome of which after transformation by a non-essentially biological procedure possesses in a stable manner a sequence of nucleotides capable of expressing a polypeptide toxic towards S.littoralis, such as that defined above. The invention also relates to the plant cells derived from their division.

The plants according to the invention are plants transformed by a non-essentially biological procedure, having in particular as predator S.littoralis, the genome of which possesses in a stable manner a sequence of nucleotides such as that defined above, capable of expressing a polypeptide toxic towards S.littoralis. The plants in question are also plants derived from their reproduction, their multiplication or hybrid crosses.

In accordance with another feature, the invention relates to plants having in particular as predator S.littoralis, possessing

in addition to their initial phenotypic and genotypic characters the property of expressing a polypeptide toxic preferentially towards S.littoralis, this property resulting from the insertion in their genome by means of genetic manipulation of a sequence of nucleotides capable of expressing the said polypeptide.

In addition, the invention relates to seeds capable of giving rise to a plant such as that defined above or derived from such a plant, characterized in that they have integrated into their genome by genetic manipulation a nucleotide sequence described above.

Other characteristics and advantages of the invention will become apparent in the course of the description and in referring to the examples in which :

- figure 1 presents the restriction map of the plasmids pHTA6 and pHTE6,

- figure 2, the restriction map of a gene for a crystal protein of the aizawai 7-29 strain cloned in the plasmid pHTA2 and defining the DNA fragments which are used as probe,

- figure 3 shows the fragment of 6.6 kb cloned in pHTA6 and the result of a hybridization carried out between this fragment and the probes described in figure 2,

- figure 4, the restriction map of the plasmid pHT671, and

- figure 5, the photographs of the immunodiffusion tests.

The hybridization experiments carried out for the implementation of the invention were performed at 42°C for 24 h in a solution containing 5 x SSC, 30% formamide and 1 Denhardt (7) in the presence of the DNA probe labelled with <sup>32</sup>P. The filters are washed at 42°C, 20 mn, by using successively the following solutions : 5 x SSC in 30% formamide, 5 x SSC, 2 x SSC, 1 x SSC and 0.5 x SSC before drying at room temperature.

EXAMPLE 1 - Construction of a DNA sequence of about 3 kb containing a hybrid gene of an insecticidal toxin.

This construction comprises :

- 1/ the preparation of gene banks of B.thuringiensis
- 2/ the selection and characterization of transforming clones containing the genes of a crystal protein and nucleotide sequences responsible

for the larvicidal activity,

3/ in vitro recombination of these sequences in a cloning vector with construction of the plasmid pHT671.

These different steps are carried out as follows :

5 1/ Preparation of gene banks of B.thuringiensis.

The total DNA of the aizawai 7-29 and entomocidus 6-01 strains of Bacillus thuringiensis is purified by using the method reported in (1) and 50 µg of each purified DNA are completely digested with the restriction enzyme PstI.

10 The DNA digested by PstI is analysed by horizontal electrophoresis on a 0.8% agarose gel and DNA fragments of a size of 5 to 8 kb are recovered from the agarose gels by electroelution in a manner described in (2).

15 The purified DNA fragments of 5-8 kb of the aizawai 7-29 strain are ligated to the DNA of the cloning vector pUC18 digested by PstI according to (3).

20 The purified DNA fragments of 5-8 kb of the entomocidus 6-01 chain are ligated to the DNA of the cloning vector pUC9 digested by PstI. The cells of E.coli JM83 are transformed with the ligation mixture as described in (4).

The transforming clones of E.coli are selected on LB medium containing 100 µg/ml of ampicillin.

25 2/ Isolation and characterization of the transforming clones containing the genes for a crystal protein.

A/ Screening of the transformed E.coli cells with the aid of an internal fragment of a gene of the crystal protein labelled with <sup>32</sup>P, used as probe :

30 Transforming clones containing recombinant plasmids carrying the gene for the crystal are detected by colony hybridization according to the method described in (5), by using as probe a PvuII fragment of 2 kb of the pBT 15-88 plasmid corresponding to an internal part of the gene for the crystal protein located on the chromosome of the berliner 1715 strain.

35 B/ Characterization of the recombinant plasmids present in the clones which react with the above probe.

Two recombinant plasmids, pHTA6 and pHTE6, isolated respectively from gene banks constructed from the strains aizawai 7-29 and entomocidus 6-01, show a homology with this probe. In each case, a DNA fragment of about 6.6 kb was cloned.

5           The restriction map of the two plasmids is given in figure 1. The comparison of the restriction sites shows that the two DNA fragments cloned appear to be identical.

10           In order to delimit the sequences corresponding to the gene for the  $\delta$ -endotoxin, different DNA fragments labelled with  $^{32}\text{P}$ , derived from a gene of the crystal previously characterized, and cloned in the recombinant plasmid pHTA2, are utilized as probes. This latter gene for the crystal also derived from the aizawai 7-29 strain codes for a protein of 130 kd active against P.brassicae but not against S.littoralis. This type of gene possesses the same restriction map  
15           as the gene for the  $\delta$ -endotoxin derived from the berliner 1715 strain. In figure 2 is shown the restriction map of this gene for the crystal protein of the aizawai 7-29 strain cloned in the plasmid pHTA2. The thick lines shown above the map correspond to the fragments used as hybridization probes.

20           The plasmids pHTA6 and pHTE6 are hydrolysed by different restriction endonucleases, analysed by horizontal electrophoresis on a 0.8% agarose gel and hybridized with the different probes.

25           The transfer of the DNA to nitrocellulose filters is carried out according to the method of SOUTHERN described in (6). The hybridization is conducted at 42°C for 24 hours in a solution containing : 5 x SSC, 30% formamide and a 1x Denhardt mixture described in (7) in the presence of a DNA probe labelled with  $^{32}\text{P}$ . The filters are then washed at 42°C for 20 minutes, by using successively the following solutions : 5 SSC in 50% formamide, 5 SSC, 2 SSC, 1 SSC  
30           and 0.5 SSC before being dried at room temperature.

35           The results of these hybridization experiments are summarized in figure 3. It appears that each extremity of the cloned DNA fragments of 6.6 kb shows a homology with the probes. The PstI-KpnI fragment of 1.5 kb reacting with the probe No. 3 corresponds to the 3' end of a gene of the crystal protein present in both the aizawai 7-29

and entomocidus 6-01 strains. As pointed out in figure 3, the probes No. 1 and No. 2 corresponding to the 5' end of the gene for the  $\delta$ -endotoxin of pHTA2 hybridize with the HindIII-HincII fragment of 1.1 kb contained in the plasmid pHTA6. The probe No. 3 which covers the 3' end of the gene of the  $\delta$ -endotoxin of pHTA2 hybridizes with the HindIII-PstI fragment of 0.4 kb contained in the plasmid pHTA6. It should be noted that a weak hybridization signal is obtained with the probe No. 2 whereas the two other probes give easily detectable signals.

From these results, the inventors have established that the HindIII-PstI DNA fragment of 3 kb corresponds to a large part of a gene for the  $\delta$ -endotoxin which commences close to the central HindIII site. It seems clear in the light of results of the hybridization experiments that the gene for the  $\delta$ -endotoxin shows substantial differences from those characterized in the prior art. On the basis of these results it was decided to clone the HindIII-PstI fragment of 3 kb in the vector pUC9.

3/ Construction of the plasmid pHT 671 containing a hybrid gene of the reconstituted insecticidal toxin.

The HindIII-HincII DNA fragment of 1.1 kb derived from the plasmid pHTe6 and the HincII-PstI DNA fragment of 1.9 kb derived from the plasmid pHTA6 are purified on agarose gels.

Equal amounts of the two purified DNA fragments and the DNA of pUC9 digested with HindIII and PstI are mixed and ligated. The ligation mixture is used to transform competent cells of E.coli JM83, then the transformed E.coli cells are selected on LB medium containing ampicillin. One of the interesting recombinant clones examined contains a plasmid designated by pHT671, the restriction map of which was determined and is shown in figure 4. This plasmid (pHT671) contains a DNA fragment of 3 kb inserted in the vector pUC9. This DNA sequence has the same restriction map as the HindIII-PstI fragments of 3 kb contained in the plasmids pHTA6 and pHTe6, but corresponds to a reconstituted DNA molecule constructed by in vitro recombination from DNA sequences derived from the aizawai 7-29 strains on the one hand and entomocidus 6-01 on the other.



EXAMPLE II : Study of the nucleotide sequence of the promoter region and of the region coding for the NH<sub>2</sub>-terminal part of the  $\delta$ -endotoxin active against the Noctuidae.

5 The HindIII-HincII fragment of pHT671 is sequenced in conformity with the method described in (8) by using a M13 system. In order to obtain partially overlapping cloned DNA fragments which will be used in the sequencing of the DNA, recourse is had to the method of subcloning by deletion in M13, developed by DALE et al (9).

10 The sequence of 940 nucleotides of the HindIII-HincII fragment which has a length of about 1 kilobase corresponds to the chain arrangement I above.

The analysis of this sequence shows that the largest open reading frame starts at position 241 and that a potential site of binding to the ribosomes, GGAGG, is found six base pairs upstream from this ATG codon (position 230 to 235). The region localized between the nucleotides 137 and 177 (position -103 to -63 upstream from the ATG codon) is strongly homologous with the region present upstream from the gene for the crystal of the strain kurstaki HD1 Dipel (BTK) sequenced by WONG et al (1983) and described in (16) and the authors of which have shown that it contains three promoters BtI, BtII, and Ec, functional in B.thuringiensis and E.coli, respectively. The comparison between the amino acid sequences deduced from the first 750 nucleotides of the genes of BTK and pHT671, show that these polypeptides exhibit significant differences at the level of the N-terminal half of the active part derived from the protoxin (only 66% strict homology). It is important to note that it is the first time that a gene for the  $\delta$ -endotoxin isolated from a strain active against the Lepidoptera codes for a polypeptide which shows substantial differences in this region. In fact, this N-terminal domain appears to be strongly conserved (more than 97% of strict homology) among all of the genes for the crystal active on Lepidoptera which have been sequenced hitherto. Moreover, the inventors have shown that the degree of variability is of the same order if the nucleotide sequences of pHT671 and other genes of the Lepidoptera type are considered.

35

EXAMPLE III : Construction of a DNA sequence of about 2.7 kb containing a gene for a larvicidal toxin.

In order to achieve this construction the DNA of the aizawai 7-29 strain of B.thuringiensis was used up to the step for the production of the plasmid pHTA6 as described in Example I.

The HindIII-PstI fragment of about 2.7 kb obtained from the plasmid pHTA6 was then subcloned in the vector pUC9, previously hydrolysed by the restriction enzymes HindIII-PstI in order to give the plasmid pHT71.

EXAMPLE IV : Study of the sequence of nucleotides constituting the plasmid pHT71 coding for a polypeptide toxic towards the larvae of Lepidoptera of the family of the Noctuidae.

The HindIII-PstI fragment of 2.7 kb of pHTA6, which was subcloned in pHT71, was sequenced by means of the technique of Sanger et al. (8) using the phage M13mp19 and the subcloning system by deletions developed by Dale et al (9). This system makes it possible to obtain M13 phages containing a series of partially overlapping DNA fragments which can be utilized for sequencing the DNA.

The sequence of nucleotides of this 2.7 kb fragment which corresponds to the chain arrangement (III) given above, was determined on the 2 DNA strands, with the exception of the last 212 nucleotides (position 2500 to 2711) which were sequenced only on a single strand.

The nucleotide sequence of this HindIII-PstI fragment has a length of 2711 nucleotides. This fragment contains the potential promoter as well as the largest part of the gene for the  $\delta$ -endotoxin active on S.littoralis.

EXAMPLE V : Study of the specific toxicity of the recombinant clones of E. coli JM83 (pHT671) and JM83 (pHT71) against S.littoralis.

The toxicity of the recombinant clones of E.coli JM83 containing pHT671 and of E.coli JM83 containing pHT71 was determined by biological tests on caterpillars of the P.brassicae and S.littoralis species as described by LECADET and MARTOURET in (10). The results were compared with the specific toxicity of the native crystal proteins purified from the strains berliner 1715 and aizawai 7-29, entomocidus 6-01 B.cereus 569 (containing the plasmid pBT45, pAMB1) against the

two species of insects. The specific toxicity of the recombinant clone and of the strains of B.thuringiensis is expressed in terms of "specificity index" previously defined.

The results obtained are reported in table 1 below.

5 In this table, for E.coli strains, the concentration 1  
corresponds to a 14 hours bacterial culture concentrated 20 times,  
disintegrated by ultrasound ; for the B.thuringiensis strains the  
concentration is expressed in  $\mu\text{g}$  of crystal protein per  $\mu\text{l}$  of  
preparation. The toxic activity of the preparations was tested by  
10 the forced ingestion with 5  $\mu\text{l}$  of preparation on caterpillars at the  
fifth stage of development, or by a technique of free ingestion  
utilizing larvae at the second stage of development.

15

20

25

30

35

TABLE 1

Comparative toxicity of a recombinant clone and two strains of B. thuringiensis towards S. littoralis and P. brassicae.

5	Strains and plasmids	S.littoralis	P.brassicae		Specificity index
		LC50 2nd larval stage	LC50 5th larval stage	LC50	LC50 S.littoralis LC50 P.brassicae
10	JM83 (pUC18)	> 1	> 1	> 1	-
	JM83 (pHT671)	0,04	0,13	0,72	0,2
	JM83 (pHTA2)	> 1	> 1	0,03	> 30
15	JM83 (pHTA4)	> 1	> 1	> 1	-
	JM83 (pHT71)	ND	0,5	> 1	< 0,5
20	berliner 1715 native crystals	ND	0,11	0,007	15,7
	aizawai 7.29 native crystals	ND	0,02	0,04	0,5
	entomocidus 601 native crystals	ND	0,028	0,012	2,3
25	B.cereus 569 (pBT45.pAM $\beta$ 1)	ND	0,38	0,054	7

30

35

Examination of the LC50 values summarized in this table 1 shows that the protein extracts of the recombinant clones JM83 (pHT671) and JM83 (pHT71) are preferentially toxic against S.littoralis. Secondly, a comparison of the values of the specificity index shows that the larvicidal activity of the recombinant clones is more specific by a factor of 2.5 times towards S.littoralis than the native crystal proteins of the aizawai strain. Moreover, the recombinant clones of JM83 (pHT671) and JM83 (pHT71) are very active against another insect of the family of the Noctuidae, Mamestra brassicae (in the case of the clone JM83 (pHT671) for example, the LC50 value is 0.02, utilizing larvae at the second stage of development).

These two results show that the gene for the larvicidal toxin constructed and cloned in the plasmids pHT671 and pHT71 codes for a protein specifically active against S.littoralis.

Other preparations obtained from recombinant clones containing plasmids carrying genes coding for other types of  $\delta$ -endotoxins (pHTA2 and pHTA4) are not active on S.littoralis : it may be seen that the plasmid pHTA2 codes for a  $\delta$ -endotoxin specifically active on P.brassicae whereas the plasmid pHTA4 codes for a  $\delta$ -endotoxin, the insect target for which has not yet been identified. It can also be seen that the crystalline inclusions produced by a strain of Bacillus cereus which has received the plasmid pBT45, one of the plasmids of the aizawai 7-29 strain which also carries a  $\delta$ -endotoxin gene (the gene of plasmid origin of the aizawai 7-29 strain), are also specifically active on P.brassicae.

Similar results are obtained by using, in the place of crude bacterial extracts, soluble protein extracts prepared from different recombinant clones of E.coli.

On the basis of the LC50 values reported in the table above and a mean individual weight of 41 mg per L5 larva (fifth larval stage) of S.littoralis, the value of the LD50 was estimated at 2.4  $\mu$ g/gram of larva for the native crystals of the aizawai 7-29 strain.

On these same bases and on the basis of equivalence factors making it possible to pass from the total bacterial mass to the

30

quantity of specific proteins (about 2% of the total proteins in E.coli JM83 (pHT671), the LD50 corresponding to the toxin produced by the expression in E.coli JM83 of the gene according to the invention cloned in the plasmid pHT671, was determined and estimated at a value close to 5.5 to 6 µg/gram of larva.

On these same bases and after determination of the LC50 of soluble protein extracts prepared from ground cultures of E.coli JM83 (pHT671), the value of the LD50 corresponding to the toxin present in these extracts was estimated at 4.15 µg/gram of larva.

In the two cases and particularly in the case of the ground preparations of E.coli, the calculated values of LD50 are approximate and higher than that of the native crystals, because it is not a question of a purified toxin. However, these data indicate without ambiguity that the gene expressed by pHT671 specifies a δ-endotoxin exhibiting the specificity towards S.littoralis. In fact, the same type of estimation made with extract of E.coli JM83 (pHTA2) carrying a δ-endotoxin gene of different specificity leads to values 30 to 50 times higher than the LD50 of the soluble extracts towards S.littoralis (135 to 350 µg/gram of larva).

The foregoing data will easily make it possible for the person skilled in the art to develop active larvicidal compositions with the proteins of the invention.

Other toxicity experiments were carried out utilizing larvae of M.brassicae, S.fruqiperda and S.littoralis at the second larval stage. The results obtained, expressed in terms of LC50 as defined for table 1, are given in table 2.

30

35

TABLE 2

ACTIVITY OF THE RECOMBINANT CLONES AGAINST  
THE LARVAE OF INSECTS OF THE FAMILY OF THE  
NOCTUIDAE: M. BRASSICAE, S. FRUGIPERDA, and  
S. LITTORALIS.

STRAINS AND PLASMIDS	INSECT LARVAE AND STAGE	<u>M. BRASSICAE</u>		<u>S. FRUGIPERDA</u>		<u>S. LITTORALIS</u>	
		LC50	2nd STAGE	LC50	2nd STAGE	LC50	2nd STAGE
JM 83 (pUC18)		NT		NT		NT	
JM 83 (pHTA2)		> 1		0,51		0,9	
JM 83 (pHT671)		0,02		0,5		0,03	
JM 83 (pHT71)		ND		ND		0,03	
JM 83 (pHTA4)		> 1		0,54		> 1	

It emerges from the examination of table 2 that the crude bacterial extracts of the recombinant clone JM83 (pHT671) are toxic towards M.brassicae and S.littoralis (the values of LC50 are 0.02 and 0.03, respectively) and weakly toxic towards S.fruqiperda (LC50 of 0.5).

The extracts of the recombinant clone E.coli JM83 (pHTA2) are weakly active towards S.fruqiperda and S.littoralis and not at all toxic towards M.brassicae. The extracts of the recombinant clone JM83 (pHTA4) are not toxic towards M.brassicae and S.littoralis and are weakly toxic toward S.fruqiperda.

These results confirm the high specific toxicity of the proteins obtained from pHT71 and pHT671 towards S.littoralis and show that this class of crystal protein is also very active towards M.brassicae.

EXAMPLE VI : Study of the specificity of the polypeptides expressed by the clones formed by introduction of the plasmids pHT671 and pHT71 into E.coli.

This study was carried out owing to immuno-diffusion tests. The results are reported in figure 5 (which includes figures 5A and 5B).

The implementation of the immuno-diffusion experiment was done in conformity with the following protocol :

Soluble extracts of proteins of E.coli clones containing the plasmids pHT671, pHTA4, pHTA2 or pHT71, pUC18 were placed in the wells Nos. 2, 3, 4, 5, 6, respectively. A sample of a solubilized purified crystal of aizawai 7-29 was placed in the well No. 1 in order to serve as positive control.

In the test recorded in figure 5A an antiserum against all of the  $\delta$ -endotoxins of aizawai 7-29, containing rabbit antibodies directed against the solubilized crystal proteins, was used and placed in the central well.

An immunoprecipitation line was observed in all of the cases except in the case of the extract of E.coli containing only the plasmid vector (well No. 6).



It was observed that the immuno-precipitation lines derived from the wells No. 4 and No. 5 cross, which shows that the products encoded by the plasmids pHTA2 and pHT71, respectively, display different antigenic determinants.

5 In the test recorded in figure 5B, the anti-serum used contained rabbit polyclonal antibodies against the crystal proteins of berliner 1715.

10 An immunoprecipitation line was observed with the extracts of E.coli JM83 (pHTA4) (well No. 3) JM83 (pHTA2) (well No. 4). On the other hand, the E.coli clones JM83 (pHT71) (well No. 5), JM83 (pHT671) (well No. 2) or JM83 (pUC9) (well No. 6) did not give immuno-precipitation.

15 It may be deduced from that that the genes for the crystal isolated in pHTA4 and pHTA2 express polypeptides having antigenic determinants in common with the proteins of the crystal of berliner 1715, a strain which is not specifically active towards S.littoralis.

20 On the other hand, the crude extracts of E.coli containing the plasmids pHT671 and pHT71 contain polypeptides having antigenic determinants in common with the crystal proteins of the aizawai 7-29 strain, which are not related immunogenically with the crystal proteins of the berliner 1715 strain.

25 These results confirm those given previously with respect to the specificity of the genes isolated in the plasmids pHT71 and pHT671.

Antigen-antibody precipitation assays have made it possible to determine the level of expression of the  $\delta$ -endotoxin genes in different recombinant clones.

30 The results obtained have shown that the crystal protein represents between 7 and 10% of the total cellular proteins of E.coli JM83 (pHTA2), between 2 and 3% in E.coli JM83 (pHT671) and between 0.5 and 1% in E.coli JM83 (pHTA4) and E.coli JM83 (pHT71).

The literature references cited in the examples  
are the following :

- 5 (1) Klier, A.F., LECADET, M-M. and DEDONDER, R., 1973,  
Sequential modifications of RNA polymerase during spo-  
rogenesis in Bacillus thuringiensis, Eur. J. Biochem., 36  
: 317-327.
- (2) MANIATIS, T., FRITSCH, E.F., SAMBROOK, J., 1982,  
Molecular cloning : A laboratory manual. Cold Spring  
10 Harbor Laboratory Press, New-York
- (3) VIEIRA, J. and MESSING, J., 1982, The pUC plasmids,  
and M13mp7 derived system for insertion mutagenesis and  
sequencing with synthetic universal primers, Gene, 19 :  
259-268.
- 15 (4) LEDERBERG, E.M. and COHEN, S.N., 1974, Transformation  
of Salmonella thyphimurium by plasmid deoxyribonucleic  
acid, J. Bacteriol., 119 : 1072-1074.
- (5) GRUNSTEIN, M. and HOGNESS, D.S., 1975, Colony hybridi-  
zation, a method for the isolation of cloned DNAs that  
20 contain a specific gene, Proc. Natl. Acad. Sci. U.S.A.,  
72 : 3961-3965.
- (6) SOUTHERN, E.M., 1975, Detection of specific sequence  
among DNA fragments separated by gel electrophoresis, J.  
Molec. Biol., 98, 503-517.
- 25 (7) DENHARDT, D.T. 1976, A membrane filter taking for the  
detection of complementary DNA. Biochem. Biophys. Res.  
Comm., 23 : 641-646.
- (8) SANGER, F., NICKLENS, S. and COULSON, A.R., 1977, DNA  
sequencing with chain terminating inhibitors. Proc. Natl.  
30 Acad. Sci. U.S.A., 74 : 5463-5467.
- (9) DALE et al. (1985) A rapid single-stranded cloning  
strategy for producing a sequential series of overlapping  
clones for use in DNA, Plasmid 1985, 13 : 31-40
- (10) LECADET, M.M. et MARTOURET D. 1987, Host specificity of  
35 the Bacillus thuringiensis  $\delta$ -endotoxin toward

Lepidopteran species : Spodoptera littoralis Bdv and  
Pieris brassicae L, J. of Invert. Pathol., 49 (n° 1) :  
37-48.

(11)CHANG et al., 1979, High frequency transformation of Bacillus subtilis protoplasts by plasmid DNA-  
Mol Gen Genet 168:111 115

(12) HEIERSON et al., 1987, Transformation of vegetative cells of Bacillus thuringiensis by plasmid DNA, Journal of Bacteriology, Mar. 1987, p. 1147-1152,

10 (13) Klier et al., 1983, Mating between Bacillus subtilis  
and Bacillus thuringiensis and transfer of cloned crystal  
genes, Mol Gen Genet (1983) 191:257 262

(14) LERECLUS et al., 1983, Isolation of a DNA, sequence related to several plasmids from Bacillus thuringiensis after a mating involving the Streptococcus faecalis plasmid pAM $\beta$ 1, Mol Gen Genet (1983) 191:307-313

(15)UMBECK et al., 1987, Genetically transformed cotton (Gossypium hirsutum L.) plants - Biotechnology vol.5 March 1987.

20 (16)WONG et al., 1983, transcriptional and translational start sites for the Bacillus thuringiensis crystal protein gene. J. of Biol. Chem., 258 : 1960-1967.

(17) OBUKOWICZ M. et al (1986). Tn<sup>5</sup> mediated integration of the  $\delta$ -endotoxin gene from B. thuringiensis into the chromosome of root colonizing Pseudomonas. J. Bacteriol., 168, 982-989.

(18) SIMON, R. et al, (1983). A broad host range mobilization system for in vivo genetic engineering : transposon mutagenesis in Gram-negative bacteria. Biotechnology, 1, pp. 784-791.

(19) Schnepf et al, (1985) The amino acid sequence of a crystal protein from Bacillus thuringiensis deduced from the DNA base sequence. J BIOL Chem 260 : 6264-6372.

(20) Adang et al, (1985) characterized full-length and truncated plasmid clones of the crystal protein of

Bacillus thuringiensis subsp. kurstaki HD-73 and their toxicity to Manduca sexta. Gene 36 : 289-300.

(21)Wabiko et al,(1986) Bacillus thuringiensis entomocidal protoxin gene sequence and gene product analysis.

5 DNA 5 : 305-314.

(22) Hofte et al, (1986) Structural and functional analysis of a cloned  $\delta$ -endotoxin gene of Bacillus thuringiensis berliner 1715. Eur J Biochem 161 : 273-280.

(23) Shibano et al, (1986) Complete structure of an insecticidal crystal protein gene from Bacillus thuringiensis. In : Bacillus molecular genetics and biotechnology applications. J.Ganesan, A.T., Hoch, J.A.(eds). Academic Press 307-320.

(24) Oeda et al, (1987) Nucleotide sequence of the insecticidal protein gene of Bacillus thuringiensis strain aizawai IPL7 and its high-level expression in Escherichia coli. Gene 53 : 113-119.